



A1(vWA1)

vWA1

&				
100				
HRP 100				
25				
TMB				



ELK Biotechnology



()



C C



1. 50 5 245 & 50
2. 100 5 495 & 100
3. 1000 5 95 & 20 5
20 245 & 50 1000
4. 100000 5 195 & 40
5 40 245 & 50 5 2000
245 & 50 100000
5. 3 100
6. PBS &



) , (

. 200





1.

2.


3.



ELK Biotechnology









●

():

1

():

3

●

vWA1

●

vWA1

2

4

8

16



/

1. 加入 100 μl 待测样品或对照品于反应孔中，轻轻混匀，室温静置 10 分钟。 2. 加入 100 μl 链霉亲和素-HRP 溶液，轻轻混匀，室温静置 10 分钟。 3. 加入 100 μl TMB 溶液，轻轻混匀，室温静置 10 分钟。 4. 加入 100 μl 终止液，轻轻混匀，室温静置 10 分钟。 5. 在 450nm 处测定吸光度 (OD 值)。		1. 加入 100 μl 待测样品或对照品于反应孔中，轻轻混匀，室温静置 10 分钟。
		2. 加入 100 μl 链霉亲和素-HRP 溶液，轻轻混匀，室温静置 10 分钟。
		3. 加入 100 μl TMB 溶液，轻轻混匀，室温静置 10 分钟。
		4. 加入 100 μl 终止液，轻轻混匀，室温静置 10 分钟。
	Avidin-HRP	5. 在 450nm 处测定吸光度 (OD 值)。
		TMB 450nmP 650nm

1. 加入 100 μl 待测样品或对照品于反应孔中，轻轻混匀，室温静置 10 分钟。

2. 加入 100 μl 链霉亲和素-HRP 溶液，轻轻混匀，室温静置 10 分钟。

3. 加入 100 μl TMB 溶液，轻轻混匀，室温静置 10 分钟。

4. 加入 100 μl 终止液，轻轻混匀，室温静置 10 分钟。



		,
		,
		30min
	,	,
	,	,
		15-30min
	,	,
		PH
	,	(
	NaN ₃ ,	NaN ₃
		,
,		TMB 450nm , 650nm



/

		,
		,
		10
		,
		,
		,
		,
		200 µL/
		,
		,
	,	,
		,
		,
		3 , 2 (2)



		3 , ,
		, , , ,
		, , , ,
	, ,	, , , ,
	, ,	6min, 3000 rpm
	, ,	, , , ,



027-59760950

elkbio@elkbiotech.cn

